

AO-A190 454

USE OF ANTI-IDIOTYPES AND SYNTHETIC PEPTIDES FOR
CONTROL OF HUMAN T-LYMPH (U) SOUTHWEST FOUNDATION FOR
BIOMEDICAL RESEARCH SAN ANTONIO TX R C KENNEDY
28 OCT 87 DAMD17-86-C-6290


1/1

UNCLASSIFIED

F/G 6/15

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A190 454

4

DTIC FILE COPY

AD

USE OF ANTI-IDIOTYPES AND SYNTHETIC PEPTIDES FOR CONTROL OF
HUMAN T-LYMPHOTROPIC VIRUS TYPE III INFECTIONS

ANNUAL PROGRESS REPORT

by

Ronald C. Kennedy, Ph.D.

October 28, 1987

Supported by

U.S. ARMY MEDICAL RESEARCH AND DEVELOPMENT COMMAND
Fort Detrick, Frederick, Maryland 21701-5012

Contract No. DAMD17-86-C-6290

SOUTHWEST FOUNDATION FOR BIOMEDICAL RESEARCH
San Antonio, Texas 78284

DOD DISTRIBUTION STATEMENT

Approved for public release; distribution unlimited.

The findings in this report are not to be construed as an official Department of the Army position unless so designated by other authorized documents.

DTIC
ELECTE
FEB 08 1988
S D

88 2 08 017

REPORT DOCUMENTATION PAGE

Form Approved
OMB No. 0704-0188

1a. REPORT SECURITY CLASSIFICATION Unclassified			1b. RESTRICTIVE MARKINGS A190454		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION/AVAILABILITY OF REPORT Approved for public release; distribution unlimited		
2b. DECLASSIFICATION/DOWNGRADING SCHEDULE					
4. PERFORMING ORGANIZATION REPORT NUMBER(S)			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Southwest Foundation for Biomedical Research		6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION		
6c. ADDRESS (City, State, and ZIP Code) P.O. Box 28147 San Antonio, TX 78284			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING/SPONSORING ORGANIZATION U.S. Army Medical Research & Develop. Command		8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER DAMD17-86-C-6290		
8c. ADDRESS (City, State, and ZIP Code) Fort Detrick Frederick, Maryland 21701-5012			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 623105	PROJECT NO. 3M2- 623105H29	TASK NO. AD
			WORK UNIT ACCESSION NO. 011		
11. TITLE (Include Security Classification) Use of Anti-idiotypes and Synthetic Peptides for Control of Human T-lymphotropic Virus Type III Infections					
12. PERSONAL AUTHOR(S) Kennedy, Ronald C.					
13a. TYPE OF REPORT Annual		13b. TIME COVERED FROM 86-09-29 TO 87-09-28		14. DATE OF REPORT (Year, Month, Day) 87/10/28	
15. PAGE COUNT 8					
16. SUPPLEMENTARY NOTATION					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP			
06	03		Synthetic peptides, anti-idiotypes, vaccines		
06	13		human immunodeficiency virus		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) During the past contract year, we have investigated the potential use of synthetic peptides and anti-idiotypes (anti-Id) for controlling HIV infection. We have identified four regions of the human immunodeficiency virus type 1 HIV-1 envelope glycoprotein that have the capacity to induce neutralizing antibody responses in experimental animals. Synthetic peptides corresponding to amino acid sequences to envelope glycoproteins gp120 and gp41 were used to identify these neutralizing epitopes. One peptide corresponding to amino acid sequences 735 to 752 from gp160 was coupled to a carrier protein and used to immunize chimpanzees. The chimpanzees produced an anti-gp160 response when immunized with this peptide; however, the neutralizing antibody response was weak. These chimpanzees, along with a control peptide immunized animal, were challenged intravenously with infectious HIV. Following challenge, the chimpanzees seroconverted and developed antibody responses against gag protein products. These studies indicated that chimpanzees were not protected from HIV infection by immunization with a					
20. DISTRIBUTION/AVAILABILITY OF ABSTRACT <input type="checkbox"/> UNCLASSIFIED/UNLIMITED <input checked="" type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION Unclassified		
22a. NAME OF RESPONSIBLE INDIVIDUAL Mrs. Virginia M. Miller			22b. TELEPHONE (Include Area Code) 301/663-7325		22c. OFFICE SYMBOL SGRD-RMI-S

single synthetic peptide that previously induced neutralizing antibodies in small experimental animals.

We also have identified two gp41 synthetic peptides that exert a profound suppression of normal human proliferative responses to mitogens and allo-antigens. Similar suppressive effects have been previously reported with a synthetic peptide analogous to amino acid sequences from the feline leukemia virus transmembrane glycoprotein.

Studies have utilized an affinity purified chimpanzee anti-gp41 as an Ab-1 preparation to generate anti-Id in rabbits. This anti-Id was serologically characterized as representing an Ab-2 gamma like preparation. BALB/c mice immunized with this anti-Id produced an Ab-3 like response which bound HIV gp41. The anti-Id induced anti-gp41 expressed a silent idiotype which was not expressed when BALB/c mice were immunized with a recombinant gp160. Thus, this anti-Id preparation can alter the serological characteristics of the immune response to gp160 in mice.

Finally, we have generated a mouse monoclonal anti-Id against a mouse monoclonal anti-CD4. This anti-Id exhibited internal image characteristics and mimicked CD4, the cellular receptor for HIV. The monoclonal anti-Id bound HIV gp120 and possessed in vitro neutralizing activity. A polyclonal anti-Id response in BALB/c mice immunized with anti-Leu3a had the capacity to neutralize four divergent HIV-1 isolates along with an HIV-2 isolate. In addition, baboons immunized with a monoclonal anti-CD4 produced an anti-Id response that recognized HIV gp120. These data suggest the vaccine possibility of an anti-Id response to a monoclonal anti-CD4 preparation.

Foreword:

In conducting the research described in the report, the investigators adhered to the "Guide for Laboratory Animal Facilities and Care," as promulgated by the Committee on the Guide for Laboratory Animal Resources, National Academy of Sciences-National Research Council.



Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

A. Immunogenicity of HIV-1 env Synthetic Peptides

Antigenic determinants are both sequential and conformational in nature. The ability to synthesize peptides predicted to contain these sequential epitopes allowed us to evaluate those domains on gp120 and gp41 which would be important in neutralization of the virus. We have identified several such regions of the HIV-1 envelope that correspond to neutralizing antigenic determinants. Peptides to be synthesized were selected by a computer model system that integrates both predicted hydrophilic and conformational information. Peptide 735-752 represented the most hydrophilic domain within gp41. Antibodies from hyperimmunized rabbits recognized gp41 as shown by immunoblotting and the env precursor (gp160) by RIP/SDS-PAGE. This result indicated that antibodies were reactive to both denatured antigen and to the more native conformational state seen by RIP analysis. Antibodies from five patients tested recognized this peptide and the specificity could be confirmed by competitive binding assays. Later studies suggested that approximately 33% of seropositive HIV-1 infected individuals recognized this peptide.

A second peptide, 503-532, consisting of the carboxyl end of gp120, was synthesized and rabbit antibodies to this peptide detected gp120 by both immunoblotting and radioimmunoprecipitation techniques. Again, antibodies from HIV-1 infected humans could detect this peptide, indicating that this peptide represents an immunogenic epitope in natural infection. Peptide 735-752 and 503-532 both elicited antibodies that neutralized the HTLV-IIIB and NY-6 strain of HIV-1, suggesting that these domains may be useful in developing HIV-1 vaccines.

We have also identified two other conserved regions within the HIV-1 envelope glycoprotein that represent neutralizing epitopes. Antisera generated against these synthetic peptides analogous to these HIV-1 envelope amino acid sequences corresponding to regions 304 to 327 and 616 to 632 also neutralized HIV-1 infectivity in vitro as assessed by reduction of reverse transcriptase activity and syncytium inhibition.

B. HIV-1 Synthetic Peptide Vaccine Studies in Chimpanzees

Peptide 735-752 analogous to gp160 amino acid sequences of HIV-1 was used to immunize two chimpanzees to evaluate if the humoral response to HIV-1 would protect this species from infection. Four immunizations of the peptide coupled to a carrier protein were performed followed by intravenous inoculation of purified cell-free virus. Prior to inoculation, antibodies from these animals were shown to react to gp41 by immunoblotting and gp160 by RIP/SDS-PAGE, indicating that humoral responses in chimpanzees were elicited. A third chimpanzee immunized with a control peptide was also challenged with virus. The humoral response was monitored on a weekly basis for the first 12 weeks following HIV infections challenge and post-inoculation reactivity to viral proteins was observed for all three primates. Antibodies to gp120, p55, and p24 were observed in all chimpanzees challenged with HIV. These data indicate that a single HIV-1 envelope glycoprotein synthetic peptide did not induce protective immunity against HIV-1 infection in chimpanzees.

C. HIV-induced Immunosuppression

We have recently demonstrated that two synthetic peptides homologous to HIV gp160 amino acid sequences 735-752 and 840-860, respectively, exert a pronounced suppression of mitogen-induced blastogenic response in vitro. The mechanism of immunosuppression remains unclear; however, our data suggest that suppression occurs at the level of IL-2 T cell interaction and that a down regulation of both IL-2 production and responsiveness may occur in HIV-peptide treated normal peripheral blood mononuclear cells. In this study, peptides conjugated to protein carriers, but not free peptides, exerted a profound suppression of the normal human lymphocyte proliferative response to ConA, PHA, PWM and alloantigens. A synthetic peptide corresponding to a 17 amino acid sequence of the HIV TAT_{III} gene product had no suppressive effects. These results suggest that, in addition to the selective cytopathic effects of HIV on CD4 bearing T-cells, viral peptide-mediated immunosuppression may also play an important role in the pathogenesis of the disease.

D. Anti-idiotypes Induce an Anti-HIV Response

We have produced in rabbits anti-idiotypic antibodies (anti-Id) against chimpanzee antibodies directed against a synthetic peptide corresponding to a native epitope associated with gp41 of HIV-1. The peptide was analogous to amino acid sequence 735-752 from gp160. Characteristics of the anti-Id preparation included: (i) detection of a shared determinant present on a second chimpanzee and one out of three rabbit antibody preparations directed against the synthetic peptide; (ii) failure to recognize an idiotypic in BALB/c mouse antisera to the peptide; and (iii) partial inhibition of the homologous chimpanzee idiotypic (Id) preparation from binding either the peptide or a recombinant HIV-1 gp160 preparation. Immunization of BALB/c mice with the anti-Id induced an anti-peptide response which bound a recombinant gp160 preparation without subsequent peptide or gp160 exposure. The anti-gp160 containing sera from mice immunized with anti-Id were used to inhibit the Id-anti-Id reaction, indicating that Id positive antibody response was induced. This Id is not normally expressed in the murine anti-gp160 immune response to the synthetic peptide and suggests that this anti-Id may activate normally silent clones. This study indicates that Id networks may be operational during the immune response to HIV-1 epitopes. In addition, non-internal image anti-Id preparations may be useful in altering the serological characteristics of an antibody response to HIV-1 relative to the nominal antigen.

E. Idiotypes Present on Monoclonal Anti-CD4 Preparations

Our laboratories have been investigating two types of molecular mimicry by anti-Id. In the first, the Id preparation (Ab-1) is an anti-CD4 and the anti-Id should mimic CD4 and bind HIV-1. In the second situation, the Id preparation is a neutralizing anti-HIV-1 and the anti-Id should mimic HIV-1 and bind CD4. The latter approach assumes that the anti-HIV-1 recognizes an HIV-1 epitope responsible for binding to CD4. We have generated a mouse monoclonal anti-Id against the mouse monoclonal anti-CD4 preparation, anti-Leu-3a. This anti-Id did not react with a panel of irrelevant mouse monoclonal antibodies, indicating that neither anti-isotype nor anti-allotype specificities were recognized. The anti-Id inhibited the ability of anti-Leu-3a to stain CD4⁺ T cells, suggesting that the anti-Id recognized an antibody combining site related determinant on anti-Leu-3a. This anti-Id bound HIV-1 determinants in

the following assays: (i) viable membrane immunofluorescence of HIV-1 infected cells; (ii) commercial ELISAs; and (iii) Western blot analysis. In addition, the anti-Id partially neutralized HIV-1 infection of T cells in vitro. These results suggest that the anti-Id reacts with an Id determinant on anti-Leu-3a and mimics part(s) of the CD4 molecule that represents the viral receptor for HIV-1.

We have also produced a polyclonal anti-Id response to anti-Leu-3a in BALB/c mice. This polyclonal anti-Id exhibited in vitro neutralizing activity against four divergent HIV-1 isolates (HTLV-IIIB, ARV-2, MN, and RF) along with an HIV-2 isolate (HIV-2_{ROD}). The anti-Id recognized anti-Leu-3a, but failed to bind another anti-human CD4 preparation (OKT4) which does not inhibit HIV binding to the CD4 molecule. In addition, the anti-Id bound gp160 in a solid phase immunoassay. Together, these results have implications for a potential AIDS vaccine utilizing anti-CD4 preparations to induce an anti-Id response with the capacity to bind HIV at its receptor site.

Two baboons were also immunized with OKT4A, a murine monoclonal antibody (MoAb) specific for the CD4 molecule on the human helper/inducer subset of T lymphocytes. Both baboons produced an anti-Id response that was specific for OKT4A. This anti-Id response also recognized other anti-CD4 MoAb, but failed to bind a series of irrelevant mouse MoAb. The anti-Id inhibited the binding of OKT4A and anti-Leu-3a to CD4⁺ cells, and stained HIV infected cells by viable membrane immunofluorescence. In addition, this anti-Id also bound HIV-1 envelope glycoproteins gp120 and gp160 by immunoblot analysis and recognized a recombinant gp160 preparation by a solid-phase immunoassay. Of particular interest, the anti-Id reacted with a band of 120 Kd in immunoblot employing simian immunodeficiency virus (SIV) antigens. Peripheral blood lymphocytes (PBL) obtained from the OKT4A-immune baboons proliferated in vitro to HIV antigens, suggesting the induction of a cell-mediated immune response. The surface phenotypes of the baboons' PBL and their in vitro proliferative response to mitogens were comparable to those of normal baboon PBL. These results suggest that anti-CD4 immunization did not result in T lymphocyte depletion or T lymphocyte anergy in these baboons. Further studies are required to assess the potential of anti-CD4 preparations as Id-based vaccines for controlling HIV infection.

Publications during the Past Year that Acknowledge Support of This Contract

Kennedy, R.C., Dreesman, G.R., Chan, T.C., Boswell, R.N., Allan, J.S., Lee, T.-H., Essex, M., Sparrow, J.T., Ho, D.D. and Kanda, P. Use of a resin bound peptide for identifying a neutralizing antigenic determinant associated with the HIV envelope. *J. Biol. Chem.* 262:5769-5774, 1987.

Chan, T.C., Dreesman, G.R. and Kennedy, R.C. Neutralization of human immunodeficiency virus by a monoclonal anti-idiotypic antibody that mimics the viral receptor. *Proc. Natl. Acad. Sci. USA* 84:3891-3895, 1987.

Dreesman, G.R., Kennedy, R.C., Kanda, P., Chan, T.C., Eichberg, J.W., Sparrow, J.T., Ho, D.D., Allan, J. S., Lee, T.H. and Essex, M. Synthetic HIV peptide vaccine candidates and protection studies in chimpanzees. In: Retroviruses of Human AIDS and Related Animal Disease (M. Girard, G. deThe and L. Valette, eds.), Marnes-la-Coquette, Paris, France, pp. 175-178, 1986.

Kennedy, R.C., Dreesman, G.R., Kanda, P., Ho, D.D., Sparrow, J.T. and Chan, T.C. Properties of synthetic peptides that identify neutralizing epitopes on the HIV envelope glycoprotein. In: Vaccines 87 (H. Ginsburg, F. Brown, R. M. Chanock, and R. A. Lerner, eds.) Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y., pp. 250-255, 1987.

Kennedy, R.C. Anti-idiotypic antibodies: Prospects in clinical and laboratory medicine. *Lab. Manag.* 25:33-41, 1987.

Kennedy, R.C., Dreesman, G.R. and Chan, T.C. Synthetic peptide and anti-idiotypic antibodies for controlling human immunodeficiency virus infection. In: International Symposium on Biotechnology in Clinical Medicine (G. Della Porta and A. Albertini, eds.) Raven Press, New York, pp. 219-225, 1987.

Kennedy, R.C., Dreesman, G.R., Lanford, R.E. and Chan, T.C. Anti-idiotypic antibodies as immunogens: Idiotypic based vaccines. In: High Technology Applications to Vaccine Development (L. Laskey and P. Berman, eds.) Blackwell Scientific, in press.

Zhou, E.M., Chan, T.C., Dreesman, G.R., Kanda, P. and Kennedy, R.C. Immune response to human immunodeficiency virus: In vivo administration of anti-idiotypic induces an anti-gp 160 response specific for a synthetic peptide, *J. Immunol.* 139:2950-2956 (1987).

Chan, T.C., Kennedy, R.C. and Kanda, P. Peptides homologous to HIV transmembrane glycoprotein suppress normal blastogenic response. *Cell. Immunol.* in press.

Kennedy, R.C. Potential applications of idiotypic based vaccines. *Vet. Immunol. Newslett.*, in press.

Dalglish, A.G., Thomson, B.T., Chan, T.C., Malkovsky, M. and Kennedy, R.C. Anti-idiotypic antibodies which mimic the T4/CD4 epitope neutralize a broad range of HIV isolates - A potential AIDS vaccine? *Lancet*, ii:1047-1050 (1987).

DISTRIBUTION LIST

12 copies	Director Walter Reed Army Institute of Research Walter Reed Army Medical Center ATTN: SGRD-UWZ-C Washington, DC 20307-5100
1 copy	Commander US Army Medical Research and Development Command ATTN: SGRD-RMI-S Fort Detrick, Frederick, Maryland 21701-5012
12 copies	Defense Technical Information Center (DTIC) ATTN: DTIC-DDAC Cameron Station Alexandria, VA 22304-6145
1 copy	Dean School of Medicine Uniformed Services University of the Health Sciences 4301 Jones Bridge Road Bethesda, MD 20814-4799
1 copy	Commandant Academy of Health Sciences, US Army ATTN: AHS-CDM Fort Sam Houston, TX 78234-6100

END

DATE

FILMED

5-88

DTIC